

~

Finite element estimates

Scientific Computing Winter 2016/2017

Lecture 19

Jürgen Fuhrmann

juergen.fuhrmann@wias-berlin.de



~

Recap

The Galerkin method

- ▶ Let V be a Hilbert space. Let $a : V \times V \rightarrow \mathbb{R}$ be a self-adjoint bilinear form, and f a linear functional on V . Assume a is coercive with coercivity constant α , and continuity constant γ .
- ▶ Continuous problem: search $u \in V$ such that

$$a(u, v) = f(v) \quad \forall v \in V$$

- ▶ Let $V_h \subset V$ be a finite dimensional subspace of V
- ▶ “Discrete” problem \equiv Galerkin approximation:
Search $u_h \in V_h$ such that

$$a(u_h, v_h) = f(v_h) \quad \forall v_h \in V_h$$

By Lax-Milgram, this problem has a unique solution as well.

Céa's lemma

- ▶ What is the connection between u and u_h ?
- ▶ Let $v_h \in V_h$ be arbitrary. Then

$$\begin{aligned}\alpha \|u - u_h\|^2 &\leq a(u - u_h, u - u_h) \quad (\text{Coercivity}) \\ &= a(u - u_h, u - v_h) + a(u - u_h, v_h - u_h) \\ &= a(u - u_h, u - v_h) \quad (\text{Galerkin Orthogonality}) \\ &\leq \gamma \|u - u_h\| \cdot \|u - v_h\| \quad (\text{Boundedness})\end{aligned}$$

- ▶ As a result

$$\|u - u_h\| \leq \frac{\gamma}{\alpha} \inf_{v_h \in V_h} \|u - v_h\|$$

- ▶ Up to a constant, the error of the Galerkin approximation is the error of the best approximation of the solution in the subspace V_h .

Definition of a Finite Element (Ciarlet)

Triplet $\{K, P, \Sigma\}$ where

- ▶ $K \subset \mathbb{R}^d$: compact, connected Lipschitz domain with non-empty interior
- ▶ P : finite dimensional vector space of functions $p : K \rightarrow \mathbb{R}^m$ (mostly, $m = 1, m = d$)
- ▶ $\Sigma = \{\sigma_1 \dots \sigma_s\} \subset \mathcal{L}(P, \mathbb{R})$: set of linear forms defined on P called *local degrees of freedom* such that the mapping

$$\begin{aligned}\Lambda_\Sigma : P &\rightarrow \mathbb{R}^s \\ p &\mapsto (\sigma_1(p) \dots \sigma_s(p))\end{aligned}$$

is bijective, i.e. Σ is a basis of $\mathcal{L}(P, \mathbb{R})$.

Local shape functions

- ▶ Due to bijectivity of Λ_Σ , for any finite element $\{K, P, \Sigma\}$, there exists a basis $\{\theta_1 \dots \theta_s\} \subset P$ such that

$$\sigma_i(\theta_j) = \delta_{ij} \quad (1 \leq i, j \leq s)$$

- ▶ Elements of such a basis are called *local shape functions*

Unisolvence

- ▶ Bijectivity of Λ_Σ is equivalent to the condition

$$\forall (\alpha_1 \dots \alpha_s) \in \mathbb{R}^s \exists! p \in P \text{ such that } \sigma_i(p) = \alpha_i \quad (1 \leq i \leq s)$$

i.e. for any given tuple of values $a = (\alpha_1 \dots \alpha_s)$ there is a unique polynomial $p \in P$ such that $\Lambda_\Sigma(p) = a$.

- ▶ Equivalent to *unisolvence*:

$$\begin{cases} \dim P = |\Sigma| = s \\ \forall p \in P : \sigma_i(p) = 0 (i = 1 \dots s) \Rightarrow p = 0 \end{cases}$$

Lagrange finite elements

- ▶ A finite element $\{K, P, \Sigma\}$ is called *Lagrange* finite element (or *nodal* finite element) if there exist a set of points $\{a_1 \dots a_s\} \subset K$ such that

$$\sigma_i(p) = p(a_i) \quad 1 \leq i \leq s$$

- ▶ $\{a_1 \dots a_s\}$: *nodes* of the finite element
- ▶ *nodal basis*: $\{\theta_1 \dots \theta_s\} \subset P$ such that

$$\theta_j(a_i) = \delta_{ij} \quad (1 \leq i, j \leq s)$$

Local interpolation operator

- ▶ Let $\{K, P, \Sigma\}$ be a finite element with shape function bases $\{\theta_1 \dots \theta_s\}$. Let $V(K)$ be a normed vector space of functions $v : K \rightarrow \mathbb{R}^m$ such that
 - ▶ $P \subset V(K)$
 - ▶ The linear forms in Σ can be extended to be defined on $V(K)$
- ▶ *local interpolation operator*

$$\mathcal{I}_K : V(K) \rightarrow P$$
$$v \mapsto \sum_{i=1}^s \sigma_i(v) \theta_i$$

- ▶ P is invariant under the action of \mathcal{I}_K , i.e. $\forall p \in P, \mathcal{I}_K(p) = p$.

Local Lagrange interpolation operator

- ▶ Let $V(K) = (C^0(K))^m$

$$\mathcal{I}_K : V(K) \rightarrow P$$

$$v \mapsto I_K v = \sum_{i=1}^s v(\mathbf{a}_i) \theta_i$$

Simplices

- ▶ Let $\{a_0 \dots a_d\} \subset \mathbb{R}^d$ such that the d vectors $a_1 - a_0 \dots a_d - a_0$ are linearly independent. Then the convex hull K of $a_0 \dots a_d$ is called *simplex*, and $a_0 \dots a_d$ are called *vertices* of the simplex.
- ▶ *Unit simplex*: $a_0 = (0 \dots 0)$, $a_1 = (0, 1 \dots 0) \dots a_d = (0 \dots 0, 1)$.

$$K = \left\{ x \in \mathbb{R}^d : x_i \geq 0 \ (i = 1 \dots d) \text{ and } \sum_{i=1}^d x_i \leq 1 \right\}$$

- ▶ A general simplex can be defined as an image of the unit simplex under some affine transformation
- ▶ F_i : face of K opposite to a_i
- ▶ \mathbf{n}_i : outward normal to F_i

Barycentric coordinates

- ▶ Let K be a simplex.
- ▶ Functions λ_i ($i = 0 \dots d$):

$$\lambda_i : \mathbb{R}^d \rightarrow \mathbb{R}$$

$$x \mapsto \lambda_i(x) = 1 - \frac{(x - a_i) \cdot \mathbf{n}_i}{(a_j - a_i) \cdot \mathbf{n}_i}$$

where a_j is any vertex of K situated in F_i .

- ▶ For $x \in K$, one has

$$\begin{aligned} 1 - \frac{(x - a_i) \cdot \mathbf{n}_i}{(a_j - a_i) \cdot \mathbf{n}_i} &= \frac{(a_j - a_i) \cdot \mathbf{n}_i - (x - a_i) \cdot \mathbf{n}_i}{(a_j - a_i) \cdot \mathbf{n}_i} \\ &= \frac{(a_j - x) \cdot \mathbf{n}_i}{(a_j - a_i) \cdot \mathbf{n}_i} = \frac{\text{dist}(x, F_i)}{\text{dist}(a_i, F_i)} \\ &= \frac{\text{dist}(x, F_i) |F_i| / d}{\text{dist}(a_i, F_i) |F_i| / d} \\ &= \frac{|K_i(x)|}{|K|} \end{aligned}$$

i.e. $\lambda_i(x)$ is the ratio of the volume of the simplex $K_i(x)$ made up of x and the vertices of F_i to the volume of K .

Barycentric coordinates II

- ▶ $\lambda_i(a_j) = \delta_{ij}$
- ▶ $\lambda_i(x) = 0 \forall x \in F_i$
- ▶ $\sum_{i=0}^d \lambda_i(x) = 1 \forall x \in \mathbb{R}^d$
(just sum up the volumes)
- ▶ $\sum_{i=0}^d \lambda_i(x)(x - a_i) = 0 \forall x \in \mathbb{R}^d$
(due to $\sum \lambda_i(x)x = x$ and $\sum \lambda_i a_i = x$ as the vector of linear coordinate functions)
- ▶ Unit simplex:
 - ▶ $\lambda_0(x) = 1 - \sum_{i=1}^d x_i$
 - ▶ $\lambda_i(x) = x_i$ for $1 \leq i \leq d$

Polynomial space \mathbb{P}_k

- ▶ Space of polynomials in $x_1 \dots x_d$ of total degree $\leq k$ with real coefficients $\alpha_{i_1 \dots i_d}$:

$$\mathbb{P}_k = \left\{ p(x) = \sum_{\substack{0 \leq i_1 \dots i_d \leq k \\ i_1 + \dots + i_d \leq k}} \alpha_{i_1 \dots i_d} x_1^{i_1} \dots x_d^{i_d} \right\}$$

- ▶ Dimension:


$$\dim \mathbb{P}_k = \binom{d+k}{k} = \begin{cases} k+1, & d=1 \\ \frac{1}{2}(k+1)(k+2), & d=2 \\ \frac{1}{6}(k+1)(k+2)(k+3), & d=3 \end{cases}$$

$$\dim \mathbb{P}_1 = d+1$$

$$\dim \mathbb{P}_2 = \begin{cases} 3, & d=1 \\ 6, & d=2 \\ 10, & d=3 \end{cases}$$





\mathbb{P}_k simplex finite elements

- ▶ K : simplex spanned by $a_0 \dots a_d$ in \mathbb{R}^d
- ▶ $P = \mathbb{P}_k$, such that $s = \dim P_k$
- ▶ For $0 \leq i_0 \dots i_d \leq k$, $i_0 + \dots + i_d = k$, let the set of nodes be defined by the points $a_{i_1 \dots i_d; k}$ with barycentric coordinates $(\frac{i_0}{k} \dots \frac{i_d}{k})$.
Define Σ by $\sigma_{i_1 \dots i_d; k}(p) = p(a_{i_1 \dots i_d; k})$.



\mathbb{P}_1 simplex finite elements








- ▶ K : simplex spanned by $a_0 \dots a_d$ in \mathbb{R}^d
- ▶ $P = \mathbb{P}_1$, such that $s = d + 1$
- ▶ Nodes \equiv vertices
- ▶ Basis functions \equiv barycentric coordinates



\mathbb{P}_2 simplex finite elements

- ▶ K : simplex spanned by $a_0 \dots a_d$ in \mathbb{R}^d
- ▶ $P = \mathbb{P}_2$, Nodes \equiv vertices + edge midpoints
- ▶ Basis functions:

$$\lambda_i(2\lambda_i - 1), (0 \leq i \leq d); \quad 4\lambda_i\lambda_j, \quad (0 \leq i < j \leq d) \quad (\text{"edge bubbles"})$$



Cuboids

- ▶ Given intervals $I_i = [c_i, d_i]$, $i = 1 \dots d$ such that $c_i < d_i$.
- ▶ *Cuboid*:

$$K = \prod_{i=1}^d [c_i, d_i]$$

- ▶ Local coordinate vector $(t_1 \dots t_d) \in [0, 1]^d$
- ▶ Unique representation of $x \in K$: $x_i = c_i + t_i(d_i - c_i)$ for $i = 1 \dots d$.
- ▶ Bijective mapping $[0, 1]^d \rightarrow K$.

Polynomial space \mathbb{Q}_k


- ▶ Space of polynomials of degree at most k in each variable
- ▶ $d = 1 \Rightarrow \mathbb{Q}_k = \mathbb{P}_k$
- ▶ $d > 1$:

$$\mathbb{Q}_k = \left\{ p(x) = \sum_{0 \leq i_1 \dots i_d \leq k} \alpha_{i_1 \dots i_d} x_1^{i_1} \dots x_d^{i_d} \right\}$$

- ▶ $\dim \mathbb{Q}_k = (k + 1)^d$

\mathbb{Q}_k cuboid finite elements

- ▶ K : cuboid spanned by intervals $[c_i, d_i]$, $i = 1 \dots d$
- ▶ $P = \mathbb{Q}_k$
- ▶ For $0 \leq i_0 \dots i_d \leq k$, let the set of nodes be defined by the points $a_{i_1 \dots i_d; k}$ with local coordinates $(\frac{i_0}{k} \dots \frac{i_d}{k})$.
Define Σ by $\sigma_{i_1 \dots i_d; k}(p) = p(a_{i_1 \dots i_d; k})$.



General finite elements

- ▶ Simplicial finite elements can be defined on triangulations of polygonal domains. During the course we will stick to this case.
- ▶ A curved domain Ω may be approximated by a polygonal domain Ω_h which is then triangulated. During the course, we will ignore this difference.
- ▶ As we have seen, more general elements are possible: cuboids, but and $T_m|_{\hat{F}} = T_n|_{\hat{F}}$ also prismatic elements etc.
- ▶ Curved geometries are possible. Isoparametric finite elements use and $T_m|_{\hat{F}} = T_n|_{\hat{F}}$ the polynomial space to define a mapping of some polyhedral reference element to an element with curved boundary

Conformal triangulations

- ▶ Let \mathcal{T}_h be a subdivision of the polygonal domain $\Omega \subset \mathbb{R}^d$ into non-intersecting compact simplices K_m , $m = 1 \dots n_e$:

$$\bar{\Omega} = \bigcup_{m=1}^{n_e} K_m$$



- ▶ Each simplex can be seen as the image of a affine transformation of a reference (e.g. unit) simplex \hat{K} :

$$K_m = T_m(\hat{K})$$

- ▶ We assume that it is conformal, i.e. if K_m, K_n have a $d - 1$ dimensional intersection $F = K_m \cap K_n$, then there is a face \hat{F} of \hat{K} and renumberings of the vertices of K_n, K_m such that $F = T_m(\hat{F}) = T_n(\hat{F})$ and $T_m|_{\hat{F}} = T_n|_{\hat{F}}$

Conformal triangulations II

- ▶ $d = 1$: Each intersection $F = K_m \cap K_n$ is either empty or a common vertex
- ▶ $d = 2$: Each intersection $F = K_m \cap K_n$ is either empty or a common vertex or a common edge



- ▶ $d = 3$: Each intersection $F = K_m \cap K_n$ is either empty or a common vertex or a common edge or a common face
- ▶ Triangulations corresponding to simplicial complexes are conformal
- ▶ Delaunay triangulations are conformal

Reference finite element

- ▶ Let $\{\widehat{P}, \widehat{K}, \widehat{\Sigma}\}$ be a fixed finite element
- ▶ Let T_K be some affine transformation and $K = T_K(\widehat{K})$
- ▶ There is a linear bijective mapping ψ_K between functions on K and functions on \widehat{K} :

$$\begin{aligned}\psi_K : V(K) &\rightarrow V(\widehat{K}) \\ f &\mapsto f \circ T_K\end{aligned}$$

- ▶ Let
 - ▶ $K = T_K(\widehat{K})$
 - ▶ $P_K = \{\psi_K^{-1}(\widehat{p}); \widehat{p} \in \widehat{P}\}$,
 - ▶ $\Sigma_K = \{\sigma_{K,i}, i = 1 \dots s : \sigma_{K,i}(p) = \widehat{\sigma}_i(\psi_K(p))\}$ Then $\{K, P_K, \Sigma_K\}$ is a finite element.

Commutativity of interpolation and reference mapping

► $\mathcal{I}_{\hat{K}} \circ \psi_K = \psi_K \circ \mathcal{I}_K,$

i.e. the following diagram is commutative:

$$\begin{array}{ccc} V(K) & \xrightarrow{\psi_K} & V(\hat{K}) \\ \downarrow \mathcal{I}_K & & \downarrow \mathcal{I}_{\hat{K}} \\ P_K & \xrightarrow{\psi_K} & P_{\hat{K}} \end{array}$$

Global interpolation operator \mathcal{I}_h

- ▶ Let $\{K, P_K, \Sigma_K\}_{K \in \mathcal{T}_h}$ be a triangulation of Ω .
- ▶ Domain:

$$D(\mathcal{I}_h) = \{v \in (L^1(\Omega))^m \text{ such that } \forall K \in \mathcal{T}_h, v|_K \in V(K)\}$$

- ▶ For all $v \in D(\mathcal{I}_h)$, define $\mathcal{I}_h v$ via

$$\mathcal{I}_h v|_K = \mathcal{I}_K(v|_K) = \sum_{i=1}^s \sigma_{K,i}(v|_K) \theta_{K,i} \quad \forall K \in \mathcal{T}_h,$$

Assuming $\theta_{K,i} = 0$ outside of K , one can write

$$\mathcal{I}_h v = \sum_{K \in \mathcal{T}_h} \sum_{i=1}^s \sigma_{K,i}(v|_K) \theta_{K,i},$$

mapping $D(\mathcal{I}_h)$ to the *approximation space*

$$W_h = \{v_h \in (L^1(\Omega))^m \text{ such that } \forall K \in \mathcal{T}_h, v_h|_K \in P_K\}$$

H^1 -Conformal approximation using Lagrangian finite elements

- ▶ Let V be a Banach space of functions on Ω . The approximation space W_h is said to be V -conformal if $W_h \subset V$.
- ▶ Non-conformal approximations are possible, we will stick to the conformal case.
- ▶ Conformal subspace of W_h with zero jumps at element faces:

$$V_h = \{v_h \in W_h : \forall n, m, K_m \cap K_n \neq \emptyset \Rightarrow (v_h|_{K_m})_{K_m \cap K_n} = (v_h|_{K_n})_{K_m \cap K_n}\}$$

- ▶ Then: $V_h \subset H^1(\Omega)$

Zero jump at interfaces with Lagrangian finite elements

- ▶ Assume geometrically conformal mesh
- ▶ Assume all faces of \widehat{K} have the same number of nodes s^∂
- ▶ For any face $F = K_1 \cap K_2$ there are renumberings of the nodes of K_1 and K_2 such that for $i = 1 \dots s^\partial$, $a_{K_1,i} = a_{K_2,i}$
- ▶ Then, $v_h|_{K_1}$ and $v_h|_{K_2}$ match at the interface $K_1 \cap K_2$ if and only if they match at the common nodes

$$v_h|_{K_1}(a_{K_1,i}) = v_h|_{K_2}(a_{K_2,i}) \quad (i = 1 \dots s^\partial)$$

Global degrees of freedom

- ▶ Let $\{a_1 \dots a_N\} = \bigcup_{K \in \mathcal{T}_h} \{a_{K,1} \dots a_{K,s}\}$
- ▶ Degree of freedom map

$$j : \mathcal{T}_h \times \{1 \dots s\} \rightarrow \{1 \dots N\}$$

$(K, m) \mapsto j(K, m)$ the global degree of freedom number

- ▶ Global shape functions $\phi_1, \dots, \phi_N \in W_h$ defined by

$$\phi_i|_K(a_{K,m}) = \begin{cases} \delta_{mn} & \text{if } \exists n \in \{1 \dots s\} : j(K, n) = i \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Global degrees of freedom $\gamma_1, \dots, \gamma_N : V_h \rightarrow \mathbb{R}$ defined by

$$\gamma_i(v_h) = v_h(a_i)$$

Lagrange finite element basis

- ▶ $\{\phi_1, \dots, \phi_N\}$ is a basis of V_h , and $\gamma_1 \dots \gamma_N$ is a basis of $\mathcal{L}(V_h, \mathbb{R})$.

Proof:


- ▶ $\{\phi_1, \dots, \phi_N\}$ are linearly independent: if $\sum_{j=1}^N \alpha_j \phi_j = 0$ then evaluation at $a_1 \dots a_N$ yields that $\alpha_1 \dots \alpha_N = 0$.
- ▶ Let $v_h \in V_h$. It is single valued in $a_1 \dots a_N$. Let $w_h = \sum_{j=1}^N v_h(a_j) \phi_j$. Then for all $K \in \mathcal{T}_h$, $v_h|_K$ and $w_h|_K$ coincide in the local nodes $a_{K,1} \dots a_{K,2}$, and by unisolvence, $v_h|_K = w_h|_K$.

Finite element approximation space


- ▶ $P_{c,h}^k = P_h^k = \{v_h \in C^0(\bar{\Omega}_h) : \forall K \in \mathcal{T}_h, v_k \circ T_K \in \mathbb{P}^k\}$
- ▶ $Q_{c,h}^k = Q_h^k = \{v_h \in C^0(\bar{\Omega}_h) : \forall K \in \mathcal{T}_h, v_k \circ T_K \in \mathbb{Q}^k\}$
- ▶ 'c' for continuity across mesh interfaces. There are also discontinuous FEM spaces which we do not consider here.

d	k	$N = \dim P_h^k$
1	1	N_v
1	2	$N_v + N_{el}$
1	3	$N_v + 2N_{el}$
2	1	N_v
2	2	$N_v + N_{ed}$
2	3	$N_v + 2N_{ed} + N_{el}$
3	1	N_v
3	2	$N_v + N_{ed}$
3	3	$N_v + 2N_{ed} + N_f$


P^1 global shape functions



P^2 global shape functions



Node based



Edge based

Global Lagrange interpolation operator

Let $V_h = P_h^k$ or $V_h = Q_h^k$

$$\mathcal{I}_h : C^0(\bar{\Omega}_h) \rightarrow V_h$$
$$v \mapsto \sum_{i=1}^N v(a_i) \phi_i$$

Further finite element constructions

- ▶ In the realm considered in this course, we stick to H^1 conformal finite elements as the weak formulations regarded work in $H^1(\Omega)$.
- ▶ With higher regularity, or for more complex problems one can construct H^2 conformal finite elements etc.
- ▶ Further possibilities for vector finite elements (divergence free etc.)

Affine transformation estimates I

- ▶ \widehat{K} : reference element
- ▶ Let $K \in \mathcal{T}_h$. Affine mapping:

$$\begin{aligned} T_K : \widehat{K} &\rightarrow K \\ \widehat{x} &\mapsto J_K \widehat{x} + b_K \end{aligned}$$

with $J_K \in \mathbb{R}^{d,d}$, $b_K \in \mathbb{R}^d$, J_K nonsingular

- ▶ Diameter of K : $h_K = \max_{x_1, x_2 \in K} \|x_1 - x_2\|$
- ▶ ρ_K diameter of largest ball that can be inscribed into K
- ▶ $\sigma_K = \frac{h_K}{\rho_K}$: local shape regularity

Affine transformation estimates II

Lemma

- ▶ $|\det J_K| = \frac{\text{meas}(K)}{\text{meas}(\widehat{K})}$
- ▶ $\|J_K\| \leq \frac{h_K}{\rho_{\widehat{K}}}$
- ▶ $\|J_K^{-1}\| \leq \frac{h_{\widehat{K}}}{\rho_K}$

Proof:

- ▶ $|\det J_K| = \frac{\text{meas}(K)}{\text{meas}(\widehat{K})}$: basic property of affine mappings
- ▶ Further:

$$\|J_K\| = \sup_{\hat{x} \neq 0} \frac{\|J_K \hat{x}\|}{\|\hat{x}\|} = \frac{1}{\rho_{\widehat{K}}} \sup_{\|\hat{x}\| = \rho_{\widehat{K}}} \|J_K \hat{x}\|$$

Set $\hat{x} = \hat{x}_1 - \hat{x}_2$ with $\hat{x}_1, \hat{x}_2 \in \widehat{K}$. Then $J_K \hat{x} = T_K \hat{x}_1 - T_K \hat{x}_2$ and one can estimate $\|J_K \hat{x}\| \leq h_K$.

- ▶ For $\|J_K^{-1}\|$ regard the inverse mapping \square

Local interpolation I

- ▶ For $w \in H^s(K)$ recall the H^s seminorm $|w|_{s,K}^2 = \sum_{|\beta|=s} \|\partial^\beta w\|_{L^2(K)}^2$

Lemma: Let $w \in H^s(K)$ and $\hat{w} = w \circ T_K$. There exists a constant c such that

$$\begin{aligned} |\hat{w}|_{s,\hat{K}} &\leq c \|J_K\|^s |\det J_K|^{-\frac{1}{2}} |w|_{s,K} \\ |w|_{s,K} &\leq c \|J_K^{-1}\|^s |\det J_K|^{\frac{1}{2}} |\hat{w}|_{s,\hat{K}} \end{aligned}$$

Proof: Let $|\alpha| = s$. By affinity and chain rule one obtains

$$\|\partial^\alpha \hat{w}\|_{L(\hat{K})} \leq c \|J_K\|^s \sum_{|\beta|=s} \|\partial^\beta w \circ T_K\|_{L^2(K)}$$

Changing variables yields

$$\|\partial^\alpha \hat{w}\|_{L(\hat{K})} \leq c \|J_K\|^s |\det J_K|^{-\frac{1}{2}} |w|_{s,K}$$

Summation over α yields the first inequality. Regarding the inverse mapping yields the second one. \square

Local interpolation II

Theorem: Let $\{\widehat{K}, \widehat{P}, \widehat{\Sigma}\}$ be a finite element with associated normed vector space $V(\widehat{K})$. Assume there exists k such that

$$\mathbb{P}_K \subset \widehat{P} \subset H^{k+1}(\widehat{K}) \subset V(\widehat{K})$$

and $H^{l+1}(\widehat{K}) \subset V(\widehat{K})$ for $0 \leq l \leq k$. There exists $c > 0$ such that for all $m = 0 \dots l+1$, $K \in \mathcal{T}_h$, $v \in H^{l+1}(K)$:

$$|v - \mathcal{I}_K^k v|_{m,K} \leq ch_K^{l+1-m} \sigma_K^m |v|_{l+1,K}$$

Draft of Proof Estimate using deeper results from functional analysis:

$$|\widehat{w} - \mathcal{I}_{\widehat{K}}^k \widehat{w}|_{m,\widehat{K}} \leq c |\widehat{w}|_{l+1,\widehat{K}}$$

(From Poincare like inequality, e.g. for $v \in H_0^1(\Omega)$, $c \|v\|_{L^2} \leq \|\nabla v\|_{L^2}$: under certain circumstances, we can estimate the norms of lower derivatives by those of the higher ones)

Local interpolation III

(Proof, continued)

Let $v \in H^{l+1}(K)$ and set $\hat{v} = v \circ T_K$. We know that $(\mathcal{I}_K^k v) \circ T_K = \mathcal{I}_{\hat{K}}^k \hat{v}$.

We have

$$\begin{aligned} |v - \mathcal{I}_K^k v|_{m,K} &\leq c \|J_K^{-1}\|^m |\det J_K|^{\frac{1}{2}} |\hat{v} - \mathcal{I}_{\hat{K}}^k \hat{v}|_{m,\hat{K}} \\ &\leq c \|J_K^{-1}\|^m |\det J_K|^{\frac{1}{2}} |\hat{v}|_{l+1,\hat{K}} \\ &\leq c \|J_K^{-1}\|^m \|J_K\|^{l+1} |v|_{l+1,K} \\ &\leq c (\|J_K\| \|J_K^{-1}\|)^m \|J_K\|^{l+1-m} |v|_{l+1,K} \\ &\leq ch_K^{l+1-m} \sigma_K^m |v|_{l+1,K} \end{aligned}$$

Local interpolation: special cases for Lagrange finite elements

- ▶ $k = 1, l = 1, m = 0$: $|v - \mathcal{I}_K^k v|_{0,K} \leq ch_K^2 |v|_{2,K}$
- ▶ $k = 1, l = 1, m = 1$: $|v - \mathcal{I}_K^k v|_{1,K} \leq ch_K \sigma_K |v|_{2,K}$

Shape regularity

- ▶ Now we discuss a family of meshes \mathcal{T}_h for $h \rightarrow 0$. We want to estimate global interpolation errors and see how they possibly diminish
- ▶ For given \mathcal{T}_h , assume that $h = \max_{K \in \mathcal{T}_h} h_j$
- ▶ A family of meshes is called *shape regular* if

$$\forall h, \forall K \in \mathcal{T}_h, \sigma_K = \frac{h_K}{\rho_K} \leq \sigma_0$$

- ▶ In 1D, $\sigma_K = 1$
- ▶ In 2D, $\sigma_K \leq \frac{2}{\sin \theta_K}$ where θ_K is the smallest angle

Global interpolation error estimate

Theorem Let Ω be polyhedral, and let \mathcal{T}_h be a shape regular family of affine meshes. Then there exists c such that for all $h, v \in H^{l+1}(\Omega)$,

$$\|v - \mathcal{I}_h^k v\|_{L^2(\Omega)} + \sum_{m=1}^{l+1} h^m \left(\sum_{K \in \mathcal{T}_h} |v - \mathcal{I}_h^k v|_{m,K}^2 \right)^{\frac{1}{2}} \leq ch^{l+1} |v|_{l+1,\Omega}$$

and

$$\lim_{h \rightarrow 0} \left(\inf_{v_h \in V_h^k} \|v - v_h\|_{L^2(\Omega)} \right) = 0$$

Global interpolation error estimate for Lagrangian finite elements, $k = 1$

- ▶ Assume $v \in H^2(\Omega)$, e.g. if problem coefficients are smooth and the domain is convex

$$\|v - \mathcal{I}_h^k v\|_{0,\Omega} + h|v - \mathcal{I}_h^k v|_{1,\Omega} \leq ch^2|v|_{2,\Omega}$$

$$|v - \mathcal{I}_h^k v|_{1,\Omega} \leq ch|v|_{2,\Omega}$$

$$\lim_{h \rightarrow 0} \left(\inf_{v_h \in V_h^k} |v - v_h|_{1,\Omega} \right) = 0$$

- ▶ If $v \in H^2(\Omega)$ cannot be guaranteed, estimates become worse. Example: L-shaped domain.
- ▶ These results immediately can be applied in Cea's lemma.

Error estimates for homogeneous Dirichlet problem

- ▶ Search $u \in H_0^1(\Omega)$ such that

$$\int_{\Omega} \lambda \nabla u \nabla v \, dx = \int_{\Omega} f v \, dx \quad \forall v \in H_0^1(\Omega)$$

Then, $\lim_{h \rightarrow 0} \|u - u_h\|_{1,\Omega} = 0$. If $u \in H^2(\Omega)$ (e.g. on convex domains) then

$$\|u - u_h\|_{1,\Omega} \leq ch|u|_{2,\Omega}$$

Under certain conditions (convex domain, smooth coefficients) one has

$$\|u - u_h\|_{0,\Omega} \leq ch|u|_{1,\Omega}$$

(“Aubin-Nitsche-Lemma”)